

## **$\beta$ -NMR: A novel spectroscopic technique in biological inorganic chemistry?**

**A. Gottberg<sup>1,2</sup>, M. Stachura<sup>1,3</sup>, M. Kowalska<sup>1</sup>, M.L. Bissell<sup>4</sup>, V. Arcisauskaite<sup>3</sup>, K. Blaum<sup>5</sup>, A. Helmke<sup>6</sup>, K. Johnston<sup>7</sup>, K. Kreim<sup>5</sup>, F.H. Larsen<sup>8</sup>, R. Neugart<sup>9</sup>, G. Neyens<sup>4</sup>, D. Szunyogh<sup>10</sup>, P.W. Thulstrup<sup>3</sup>, D.T. Yordanov<sup>5</sup>, L. Hemmingsen<sup>3</sup>**

<sup>1</sup> CERN, Geneva, Switzerland. Monika.stachura@cern.ch

<sup>2</sup> Instituto de Estructura de la Materia, CSIC, Serrano 113bis, 28006-Madrid, Spain.

<sup>3</sup> Kemisk Institut, Københavns Universitet, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark. lhe@chem.ku.dk

<sup>4</sup> Instituut voor Kern- en Stralingsfysica, KU Leuven, Celestijnenlaan 200 D, B-3001 Leuven, Belgium.

<sup>5</sup> Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany.

<sup>6</sup> Humedics GmbH, Marie-Elisabeth-Lüders-Str. 1, 10625 Berlin, Germany.

<sup>7</sup> Universität des Saarlandes, Experimentalphysik, 66123 Saarbrücken, Germany.

<sup>8</sup> Institut for Fødevarevidenskab, Københavns Universitet, Rolighedsvej 30, 1958 Frederiksberg C, Denmark.

<sup>9</sup> Institut für Kernchemie, Universität Mainz, Fritz-Straßmann-Weg 2, 55128 Mainz, Germany.

<sup>10</sup> Szervetlen és Analitikai Kémiai Tanszék, Szegedi Tudományegyetem, Dom ter 7, 6720 Hungary.

Nuclear magnetic resonance (NMR) spectroscopy is a powerful and versatile technique, which allows for elucidation of protein structure and dynamics. However, several biologically relevant metal ions, such as  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cu}^+$  or  $\text{Zn}^{2+}$  are problematic in conventional NMR spectroscopy due to the lack of suitable stable isotopes and limited sensitivity. Moreover, many other analytical tools fail for the same metal ions because of the closed electronic shell structure. Here we present an experimental NMR approach that is one billion times more sensitive than conventional NMR spectroscopy. The increase in sensitivity is achieved by recording the anisotropic emission of  $\beta$ -particles in the decay of highly spin-polarized nuclei<sup>1,2</sup>.  $\beta$ -NMR may be applied to many chemical elements and we aim to study e.g.  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cu}^+$  and  $\text{Zn}^{2+}$ . The technique has been applied in solid-state physics<sup>3,4</sup> but not yet in biological inorganic chemistry. Although there are still technological challenges to overcome,  $\beta$ -NMR spectroscopy holds considerable promise as a novel very sensitive technique in biological inorganic chemistry.

## **References:**

1. Matthias E, Olsen B, Shirley DA, Templeton JE, Steffen RM (1971) Phys. Rev. A4:1626-1658
2. Gottberg A, Stachura M, Kowalska M, Bissell M.L, Arcisauskaite V, Blaum K, Helmke A, Johnston K, Kreim K, Larsen FH, Neugart R, Neyens G, Szunyogh D, Thulstrup PW, Yordanov DT, Hemmingsen L (2014) *manuscript in preparation*
3. Mansour AI, Morris GD, Salman Z, Chow KH, Dunlop T, Jung J, Fan I, MacFarlane WA, Kiefl RF, Parolin TJ, Saadaoui H, Wang D, Hossain MD, Song Q, Smadella M, Mosendz O, Kardasz B, Heinrich B (2007) Physica B 401-402:662-665
4. Salman Z, Wang D, Chow KH, Hossain MD, Kreitzman SR, Keeler TA, Levy CDP, MacFarlane WA, Miller RI, Morris GD, Parolin TJ, Saadaoui H, Smadella M, Kiefl RF (2007) Phys. Rev. Lett. 98:167001

